arXiv:1603.01818v2 [math.AP] 22 Dec 2016

Well-posedness of a porous medium flow with fractional pressure in Sobolev spaces

Xuhuan Zhou[1] Weiliang Xiao[2*]

Abstract. The nonnegative solution for a linear degenerate diffusion transport eqution is proved. As a result, we show the existence and uniqueness of the solution for the fractional porous medium equation in Sobolev spaces H^α with nonnegative initial data, $\alpha > \frac{d}{2} + 1$. Besides, we correct a mistake in our previous paper [13].

§1 Introduction

In this paper we consider the following porous medium type equation

$$\partial_t u = \nabla \cdot (u \nabla p), \quad p = (-\Delta)^{-s} u, \quad 0 < s < 1. \tag{1.1}$$

Where $x \in \mathbb{R}^n$, $n \geq 2$, and $t > 0$. The initial data $u(x, 0) \geq 0$.

This model is based on Darcy's law and the pressure is given by an inverse fractional Laplacian operator. It was first introduced by Caffarelli and Vázquez [4], in which they proved the existence of a weak solution when u_0 is a bounded function with exponential decay at infinity. For $\alpha = \frac{n}{n+2-2s}$, Caffarellin, Soria and Vázquez [3] proved that the bounded nonnegative solutions are C^α continuous in a strip of space-time for $s \neq 1/2$. And same conclusion for the index $s = 1/2$ was proved by Caffarelli and Vázquez in [5]. [6, 7, 12] give a detail description of the large-time asymptotic behaviour of the solutions of (1.1). [2,11] considered some degenerate cases and show the existence and properties of self-similar solutions. Allen, Caffarelli and Vasseur [1] studied the equation with another fractional time derivative, and proved the Hölder continuity for its weak solutions.

In this paper, we study the existence and uniqueness of solutions of (1.1) in Sobolev spaces. The method we used here is novel: Unlike the usual way to consider the weak solution in L^∞ or construct approximate solutions of linear transport systems, we solve equation (1.1) by constructing linear degenerate diffusion transport systems. The well-posedness and properties of the constructed linear degenerate diffusion transport are interesting problem themselves. By this way we get that for $s \in [\frac{1}{2}, 1)$, $\alpha > \frac{d}{2} + 1$, $u_0 \in H^\alpha(\mathbb{R}^n)$ nonnegative, then for some $T_0 > 0$, the unique solution of (1.1) in $\mathbb{R}^n \times [0, T_0]$ exists. Besides, using the methods and results in this paper, we correct a mistake in our previous paper [13].

Mathematics Subject Classification(2000): 35K55, 35K65, 76S05.
Keywords: fractional porous medium equation, degenerate diffusion transport equation, Sobolev spaces.

§2 Preliminaries

Define $\rho(x) \in C_c^\infty(\mathbb{R}^n)$ by

$$\rho(x) = \begin{cases} c_0 \exp(-\frac{1}{1-|x|^2}), |x| < 1, \\ 0, |x| \geq 1, \end{cases}$$

where c_0 is selected such that $\int \rho(x)dx = 1$. The operator J_ϵ is defined by

$$J_\epsilon u = \rho_\epsilon * u = \epsilon^{-n}\rho(\frac{\cdot}{\epsilon}) * u,$$

and it has the following properties:

Proposition 2.1. *(1)* $\Lambda^s J_\epsilon u = J_\epsilon \Lambda^s u$, $s \in \mathbb{R}$.
(2) For all $u \in L^p(\mathbb{R}^n)$, $v \in H^\alpha(\mathbb{R}^n)$, *with* $\frac{1}{p} + \frac{1}{q} = 1$, $\int (J_\epsilon f)g = \int f(J_\epsilon g)$.
(3) For all $u \in H^\alpha(\mathbb{R}^n)$,

$$\lim_{\epsilon \to 0} \|J_\epsilon u - u\|_{H^\alpha} = 0, \quad \lim_{\epsilon \to 0} \|J_\epsilon u - u\|_{H^{\alpha-1}} \leq C\|u\|_{H^\alpha}.$$

(4) For all $u \in H^\alpha(\mathbb{R}^n)$, $s \in \mathbb{R}$, $k \in \mathbb{Z} \cup \{0\}$, *then*

$$\|J_\epsilon u\|_{H^{\alpha+k}} \leq \frac{C_{\alpha k}}{\epsilon^k}\|u\|_{H^\alpha}, \quad \|J_\epsilon D^k u\|_{L^\infty} \leq \frac{C_k}{\epsilon^{\frac{n}{2}+k}}\|u\|_{H^\alpha}$$

Following propositions can be found in [8, 9].

Proposition 2.2. *Suppose that* $s > 0$ *and* $1 < p < \infty$. *If* $f, g \in \mathcal{S}$, *the Schwartz class, then we have*

$$\|\Lambda^s(fg) - f\Lambda^s g\|_{L^p} \leq c\|\nabla f\|_{L^{p_1}}\|g\|_{\dot{H}^{s-1,p_2}} + c\|g\|_{L^{p_4}}\|f\|_{\dot{H}^{s,p_3}}$$

and

$$\|\Lambda^s(fg)\|_{L^p} \leq c\|f\|_{L^{p_1}}\|g\|_{\dot{H}^{s,p_2}} + c\|g\|_{L^{p_4}}\|f\|_{\dot{H}^{s,p_3}}$$

with $p_2, p_3 \in (1, +\infty)$ *such that* $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p_3} + \frac{1}{p_4}$.

Proposition 2.3. *Let* $0 \leq s \leq 2$, $f \in \mathcal{S}(\mathbb{R}^n)$, *we have the pointwise inequality,*

$$2f(x)\Lambda^s f(x) \geq \Lambda^s f^2(x).$$

Proposition 2.4. *Let* α_1 *and* α_2 *be two real numbers such that* $\alpha_1 < \frac{n}{2}$, $\alpha_2 < \frac{n}{2}$ *and* $\alpha_1 + \alpha_2 > 0$. *Then there exists a const* $C = C_{\alpha_1,\alpha_2} \geq 0$ *such that for all* $f \in \dot{H}^{\alpha_1}$ *and* $g \in \dot{H}^{\alpha_2}$,

$$\|fg\|_{\dot{H}^\alpha} \leq C\|f\|_{\dot{H}^{\alpha_1}}\|g\|_{\dot{H}^{\alpha_2}},$$

where $\alpha = \alpha_1 + \alpha_2 - \frac{n}{2}$.

§3 Main Results

Theorem 3.1. *Let* $s \in [\frac{1}{2}, 1]$, $T > 0$, $\alpha > \frac{n}{2} + 1$, $u_0 \in H^\alpha(\mathbb{R}^n)$, $v \in C([0,T]; H^\alpha(\mathbb{R}^n))$, *and* $v \geq 0$. *Then there is a unique solution* $u \in C^1([0,T]; H^\alpha(\mathbb{R}^n))$ *to the linear initial value problem*

$$\begin{cases} \partial_t u = \nabla u \cdot \nabla(-\Delta)^{-s} v - v(-\Delta)^{1-s} u, \\ u(x,0) = u_0. \end{cases} \tag{3.1}$$

And if the initial data $u_0 \geq 0$, *we can get* $u \geq 0, (x,t) \in \mathbb{R}^n \times [0,T]$.

Proof. For any $\epsilon > 0$, we consider the following linear problem

$$\begin{cases} \partial_t u^\epsilon = F_\epsilon(u^\epsilon) = J_\epsilon(\nabla J_\epsilon u^\epsilon \cdot \nabla(-\Delta)^{-s} v) - J_\epsilon(v(-\Delta)^{1-s} J_\epsilon u^\epsilon), \\ u^\epsilon(x,0) = u_0. \end{cases} \tag{3.2}$$

By Proposition 2.1, Proposition 2.2 and $s \geq \frac{1}{2}$ we can estimate

$$\begin{aligned} \|F_\epsilon(u_1^\epsilon) - F_\epsilon(u_2^\epsilon)\|_{H^\alpha} &= \|J_\epsilon(\nabla J_\epsilon(u_1^\epsilon - u_2^\epsilon) \cdot \nabla(-\Delta)^{-s} v) - J_\epsilon(v(-\Delta)^{1-s} J_\epsilon(u_1^\epsilon - u_2^\epsilon)\|_{H^\alpha} \\ &\leq C(\epsilon, \|v\|_{H^\alpha}) \|u_1^\epsilon - u_2^\epsilon\|_{H^\alpha}. \end{aligned}$$

By Picard Theorem, for any $\alpha > \frac{n}{2} + 1, \epsilon > 0$, there exists a $T_\epsilon = T_\epsilon(u_*) > 0$, problem (3.2) has a unique solution $u^\epsilon \in C^1([0, T_\epsilon); H^\alpha)$. By Proposition 2.1 and Proposition 2.3,

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|u^\epsilon\|_{L^2}^2 &= \int \nabla J_\epsilon u^\epsilon \cdot \nabla(-\Delta)^{-s} v J_\epsilon u^\epsilon - \int v(-\Delta)^{1-s} J_\epsilon u^\epsilon J_\epsilon u^\epsilon \\ &\leq \frac{1}{2}\int \nabla |J_\epsilon u^\epsilon|^2 \cdot \nabla(-\Delta)^{-s} v - \frac{1}{2}\int v(-\Delta)^{1-s} |J_\epsilon u^\epsilon|^2 \\ &\leq \frac{1}{2}\int |J_\epsilon u^\epsilon|^2 (-\Delta)^{1-s} v - \frac{1}{2}\int |J_\epsilon u^\epsilon|^2 (-\Delta)^{1-s} v = 0. \end{aligned}$$

Moreover, for any $\alpha > 0$,

$$\begin{aligned} \frac{1}{2}\frac{d}{dt}\|\Lambda^\alpha u^\epsilon\|_{L^2}^2 &= \int \Lambda^\alpha(\nabla J_\epsilon u^\epsilon \cdot \nabla(-\Delta)^{-s} v) J_\epsilon \Lambda^\alpha u^\epsilon - \int \Lambda^\alpha(v(-\Delta)^{1-s} J_\epsilon u^\epsilon) \Lambda^\alpha J_\epsilon u^\epsilon \\ &\leq C\|[\Lambda^\alpha, \nabla(-\Delta)^{-s} v] \nabla J_\epsilon u^\epsilon\|_{L^2} \|\Lambda^\alpha u^\epsilon\|_{L^2} + \int \nabla(-\Delta)^{-s} v \Lambda^\alpha \nabla J_\epsilon u^\epsilon \Lambda^\alpha J_\epsilon u^\epsilon \\ &+ C\|[\Lambda^\alpha, v](-\Delta)^{1-s} J_\epsilon u^\epsilon\|_{L^2} \|\Lambda^\alpha u^\epsilon\|_{L^2} - \int v \Lambda^\alpha (-\Delta)^{1-s} J_\epsilon u^\epsilon \Lambda^\alpha J_\epsilon u^\epsilon \end{aligned}$$

By Proposition 2.2 and Sobolev embedding,

$$\begin{aligned} \|[\Lambda^\alpha, \nabla(-\Delta)^{-s} v] \nabla J_\epsilon u^\epsilon\|_{L^2} &\leq C\|(-\Delta)^{1-s} v\|_{L^\infty} \|\Lambda^{\alpha-1} \nabla J_\epsilon u^\epsilon\|_{L^2} + \|\nabla(-\Delta)^{-s} v\|_{\dot{H}^\alpha} \|\nabla J_\epsilon u^\epsilon\|_{L^\infty} \\ &\leq C\|v\|_{H^\alpha} \|u^\epsilon\|_{H^\alpha}, \\ \|[\Lambda^\alpha, v](-\Delta)^{1-s} J_\epsilon u^\epsilon\|_{L^2} &\leq C\|\nabla v\|_{L^\infty} \|(-\Delta)^{1-s} J_\epsilon u^\epsilon\|_{\dot{H}^{\alpha-1}} + \|v\|_{\dot{H}^\alpha} \|(-\Delta)^{1-s} J_\epsilon u^\epsilon\|_{L^\infty} \\ &\leq C\|v\|_{H^\alpha} \|u^\epsilon\|_{H^\alpha}. \end{aligned}$$

By Proposition 2.3,

$$\begin{aligned} &\int \nabla(-\Delta)^{-s} v \Lambda^\alpha \nabla J_\epsilon u^\epsilon \Lambda^\alpha J_\epsilon u^\epsilon - \int v \Lambda^\alpha (-\Delta)^{1-s} J_\epsilon u^\epsilon \Lambda^\alpha J_\epsilon u^\epsilon \\ &\leq \frac{1}{2}\int \nabla(-\Delta)^{-s} v \nabla(\Lambda^\alpha J_\epsilon u^\epsilon)^2 - \frac{1}{2}\int v(-\Delta)^{1-s} (\Lambda^\alpha J_\epsilon u^\epsilon)^2 \\ &\leq C\|v\|_{H^\alpha} \|u^\epsilon\|_{H^\alpha}^2. \end{aligned}$$

Combine the above estimates,

$$\frac{d}{dt}\|u^\epsilon(\cdot, t)\|_{H^\alpha} \leq C\|v\|_{H^\alpha} \|u^\epsilon\|_{H^\alpha}.$$

By Gronwall's inequality,

$$\|u^\epsilon(\cdot, t)\|_{H^\alpha} \leq \|u_0\|_{H^\alpha} \exp(C \sup_{0 \leq t \leq T} \|v\|_{H^\alpha}).$$

Such the solution u^ϵ exists on $[0, T]$. Similarly,

$$\frac{d}{dt}\|u^\epsilon(\cdot, t)\|_{H^{\alpha-1}} \leq C\|v\|_{H^\alpha} \|u^\epsilon\|_{H^\alpha} \leq C(\|v\|_{H^\alpha}, \|u_0\|_{H^\alpha}, T).$$

By Aubin compactness theorem, there is a subsequence of $\{u^{\frac{1}{n}}\}_{n\geq 1}$ that convergence strongly to u in $C([0,T];H^\alpha)$. If $\alpha > \frac{d}{2}+1$, $H^\alpha \hookrightarrow C^1$, so u is a solution of (3.1).

If $u, \tilde{u}$ are two solutions of problem (3.1), then $w = u - \tilde{u}$ satisfies

$$\begin{cases} \partial_t w = \nabla w \cdot \nabla(-\Delta)^{-s} v - v(-\Delta)^{1-s} w, \\ w(x,0) = 0. \end{cases}$$

Similarly, we can get $\frac{d}{dt}\|w\|_{L^2} \leq 0$ and $\frac{d}{dt}\|w\|_{\dot{H}^\alpha} \leq \|v\|_{H^\alpha}\|w\|_{H^\alpha}$, i.e, $\frac{d}{dt}\|w\|_{H^\alpha} \leq \|v\|_{H^\alpha}\|w\|_{H^\alpha}$. By Gronwall's inequality we can deduce $u(x,t) = 0$, $(x,t) \in \mathbb{R}^n \times [0,T]$.

Since $u_0 \geq 0$ then if there exists a first time t_0 where for some point x_0 we have $u(x_0,t_0) = 0$, then (x_0,t_0) will correspond to a minimum point and therefore $\nabla u(x_0,t_0) = 0$, and

$$(-\Delta)^{1-s}u(x) = c\int \frac{u(x)-u(y)}{|y|^{n+2-2s}}dy \leq 0.$$

Hence $u_t|_{(x_0,t_0)} \geq 0$. So $u(x,t) \geq 0$ for all $(x,t) \in \mathbb{R}^n \times [0,T]$. □

Theorem 3.2. *Let $n \geq 2$, $s \in [\frac{1}{2}, 1)$, $\alpha > \frac{d}{2}+1$, $u_0 \in H^\alpha(\mathbb{R}^n)$, and $u_0 \geq 0$. Then there is a unique solution $u \in C^1([0,T_0], H^\alpha(\mathbb{R}^n))$ to the linear initial value problem*

$$\begin{cases} \partial_t u = \nabla \cdot (u\nabla(-\Delta)^{-s}u), \\ u(x,0) = u_0. \end{cases}$$

And if the initial data $u_0 \geq 0$, we can get $u \geq 0, (x,t) \in \mathbb{R}^n \times [0,T_0]$.

Proof. Set $u^1 = u_0$. Notice $\partial_t u = \nabla \cdot (u\nabla(-\Delta)^{-s}u) = \nabla u \cdot \nabla(-\Delta)^{-s}u - u(-\Delta)^{1-s}u$, we construct a sequence $\{u^n\}$ defined by solving the following systems

$$\begin{cases} \partial_t u^{n+1} = \nabla u^{n+1} \cdot \nabla(-\Delta)^{-s}u^n - u^n(-\Delta)^{1-s}u^{(n+1)}, \\ u^{n+1}(x,0) = u_0. \end{cases} \tag{3.3}$$

Firstly by Theorem 3.1, we get $u^2 \in C([0,T);H^\alpha)$, $\forall T < \infty$, and it satisfies $u^2 \geq 0$ and

$$\sup_{0\leq t\leq T}\|u^2\|_{H^\alpha} \leq \|u_0\|_{H^\alpha}\exp(C\|u^1\|_{H^\alpha}T).$$

If $\exp(2C\|u_1\|_{H^\alpha}T_0) \leq 2$, for example $T_0 = \frac{\ln 2}{2C(1+\|u_0\|_{H^\alpha})}$, we have $\sup_{0\leq t\leq T_0}\|u^2\|_{H^\alpha} \leq 2\|u_0\|_{H^\alpha}$. By the standard induction argument, if $u^n \in C([0,T_0];H^\alpha)$, $u^n \geq 0$ is a solution of (3.3) with $\|u^n\|_{H^\alpha} \leq 2\|u_0\|_{H^\alpha}$, by Theorem 3.1 we can get $u^{n+1} \in C([0,T_0];H^\alpha)$, $u^{n+1} \geq 0$ and

$$\sup_{0\leq t\leq T_0}\|u^{n+1}\|_{H^\alpha} \leq \|u_0\|_{H^\alpha}\exp(C\|u^n\|_{H^\alpha}T_0) \leq 2\|u_0\|_{H^\alpha},$$
$$\frac{d}{dt}\|u^{n+1}\|_{H^{\alpha-1}} \leq C\|u^n\|_{H^\alpha}\|u^{n+1}\|_{H^\alpha} \leq C\|u_0\|^2_{H^\alpha}.$$

By Aubin compactness theorem, there is a subsequence of u^n that convergence strongly to u in $C([0,T];H^\alpha)$. If $u \geq 0, \tilde{u} \geq 0$ are two solutions of problem (3.1), then $w = u - \tilde{u}$ satisfies that

$$\begin{cases} \partial_t w = \nabla \cdot (w\nabla(-\Delta)^{-s}u) + \nabla \cdot (\tilde{u}\nabla(-\Delta)^{-s}w), \\ w(x,0) = 0. \end{cases}$$

By Proposition 2.2 we can get

$$\frac{1}{2}\frac{d}{dt}\|w\|^2_{L^2} = \int w\nabla \cdot (w\nabla(-\Delta)^{-s}u) + \int w\nabla\tilde{u} \cdot \nabla(-\Delta)^{-s}w - \int w\tilde{u}(-\Delta)^{1-s}w \triangleq I_1 + I_2 + I_3.$$

I_1, I_3 can be estimated as:

$$I_1 = \int w\nabla w \cdot \nabla(-\Delta)^{-s}u = \frac{1}{2}\int \nabla w^2 \cdot \nabla(-\Delta)^{-s}u = \frac{1}{2}\int w^2(-\Delta)^{1-s}u \le C\|u\|_{H^\alpha}\|w\|_{L^2}^2,$$

$$I_3 \le -\frac{1}{2}\int \tilde{u}(-\Delta)^{1-s}w^2 = \int -\frac{1}{2}(-\Delta)^{1-s}\tilde{u}\cdot w^2 \le C\|u\|_{H^\alpha}\|w\|_{L^2}^2.$$

When $s > \frac{1}{2}$,

$$\begin{aligned} I_2 &\le C\|w\|_{L^2}\|\nabla u \cdot \nabla(-\Delta)^{-s}w\|_{L^2} \\ &\le C\|w\|_{L^2}\|\nabla u\|_{\dot{H}^{\frac{n}{2}+1-2s}}\|\nabla(-\Delta)^{-s}w\|_{\dot{H}^{2s-1}} \le C\|u\|_{H^\alpha}\|w\|_{L^2}^2. \end{aligned}$$

When $s = \frac{1}{2}$, the above estimates are still valid. Combine the above estimates, $\frac{d}{dt}\|w\|_{L^2} \le C\|w\|_{L^2}\|u\|_{H^\alpha}$. By Gronwall's inequality we can deduce $w(x,t) = 0$ on $[0, T_0]$. □

§4 Correction

In our previous paper [13] in which we establish the well-posedness result in Besov spaces for the equation (1.1), there is a mistake in page 9 when we estimate the term J_4' in equation (4.5). To correct the mistake, we modify our proof in the way as following: First we construct the approximate equation:

$$\begin{cases} u_t^{(n+1)} = \nabla u^{(n+1)} \cdot \nabla(-\triangle)^{-s}u_\epsilon^{(n)} - u_\epsilon^{(n)}(-\triangle)^{1-s}u^{(n+1)}; \\ u^{(n+1)}(0) = \sigma_\epsilon * u_0, \quad u^{(1)} = \sigma_\epsilon * u_0. \end{cases} \tag{4.1}$$

By the argument in section 2, we can always find the sequence $u^{(n)}$ who solves the linear systems (4.1). Assume $u_0 \ge 0$, we prove $u^{(n+1)} \ge 0$. Inspired by [4], we assume that x_0 is a point of minimum of $u^{(n+1)}$ at time $t = t_0$. This indicate that $\nabla u^{(n+1)}(x_0) = 0$, and

$$(-\triangle)^{1-s}u^{(n+1)}(x_0) = c\int \frac{u(x_0) - u(y)}{|y|^{n+2(1-s)}}dy \le 0.$$

Thus we deduce $\frac{\partial}{\partial t}u^{(n+1)}\big|_{t=t_0} \ge 0$, and by induction there holds

$$u^{(n+1)} \ge 0.$$

By the same way as [13], taking $\triangle_j$ on (4.1), we obtain

$$\begin{aligned} \partial_t \triangle_j u^{(n+1)} = &\sum [\triangle_j, \partial_i(-\triangle)^{-s}u_\epsilon^{(n)}]\partial_i u^{n+1} + \sum \partial_i(-\triangle)^{-s}u_\epsilon^{(n)}\triangle_j(\partial_i u^{(n+1)}) \\ &- [\triangle_j, u_\epsilon^{(n)}](-\triangle)^{1-s}u^{(n+1)} - u_\epsilon^{(n)}\triangle_j(-\triangle)^{1-s}u^{(n+1)}. \end{aligned}$$

Multiplying both sides by $\frac{\triangle_j u^{(n+1)}}{|\triangle_j u^{(n+1)}|}$, and integrating over $\mathbb{R}^d$, then denote the corresponding part in the right side by J_1', J_2', J_3', J_4', respectively. We obtain the estimates,

$$J_1' \le C2^{-j\alpha}\|u^{(n+1)}\|_{B^\alpha_{1,\infty}}\|u^{(n)}\|_{B^{\alpha+1-2s}_{1,\infty}}$$

$$J_2' \le C2^{-j\alpha}\|u^{(n+1)}\|_{B^\alpha_{1,\infty}}\|u^{(n)}\|_{B^{\alpha+1-2s}_{1,\infty}}$$

$$J_3' \le C2^{-j\alpha}\|u^{(n)}\|_{B^\alpha_{1,\infty}}\|u^{(n+1)}\|_{B^{\alpha+1-2s}_{1,\infty}}.$$

And the estimate for the term J_4' is replaced by

$$\begin{aligned} J_4' &= -\int u^{(n)} \triangle_j (-\triangle)^{1-s} u^{(n+1)} \frac{\triangle_j u^{(n+1)}}{|\triangle_j u^{(n+1)}|} \\ &\leq -\int u^{(n)} (-\triangle)^{1-s} |\triangle_j u^{(n+1)}| \\ &\leq -\int (-\triangle)^{1-s} u^{(n)} |\triangle u^{(n+1)}| \\ &\leq 2^{-j\alpha} \|u^n\|_{B^{r+2-2s}_{1,\infty}} \|u^{(n+1)}\|_{B^{\alpha}_{1,\infty}}. \end{aligned}$$

Here $r > d$ be a arbitrary real number, and in the first inequality we use the following pointwise estimate:

Proposition 4.1. *[10] Set $0 \leq \alpha \leq 2$, $p \geq 1$. Then for any $f \in \mathcal{S}(\mathbb{R}^d)$, there holds*

$$p|f(x)|^{p-2} f(x) \Lambda^{\alpha} f(x) \geq \Lambda^{\alpha} |f(x)|^p.$$

Taking r such that $r + 2 - 2s < \alpha$, e.g. set $r = \alpha - 1$, we conclude

$$\frac{d}{dt} \|u^{(n+1)}\|_{B^{\alpha}_{1,,\infty}} \leq \|u^{(n)}\|_{B^{\alpha}_{1,,\infty}} \|u^{(n+1)}\|_{B^{\alpha}_{1,,\infty}}.$$

The other parts of the proof are no difference.

Acknowledgement

We are very grateful for PhD. Mitia Duerinckx for pointing out our mistake and share some good views on this problem. This paper is supported by the NNSF of China under grants No. 11601223 and No.11626213.

References

[1] M. Allen, L. Caffarelli, A. Vasseur. *Porous medium flow with both a fractional potential pressure and fractional time derivative.* arXiv preprint arXiv:1509.06325, 2015.

[2] P. Biler, C. Imbert, G. Karch. *The nonlocal porous medium equation: Barenblatt profiles and other weak solutions.* Archive for Rational Mechanics and Analysis, 2015, 215(2): 497-529.

[3] L. Caffarelli, F. Soria, J. L. Vázquez. *Regularity of solutions of the fractional porous medium flow.* arXiv preprint arXiv:1201.6048, 2012.

[4] L. Caffarelli, J. L. Vázquez. *Nonlinear porous medium flow with fractional potential pressure.* Archive for rational mechanics and analysis, 2011, 202(2): 537-565.

[5] L. Caffarelli, J. L. Vázquez. *Regularity of solutions of the fractional porous medium flow with exponent* 1/2. arXiv preprint arXiv:1409.8190, 2014.

[6] L. Caffarelli, J. L. Vázquez. *Asymptotic behaviour of a porous medium equation with fractional diffusion.* arXiv preprint arXiv:1004.1096, 2010.

[7] J. A. Carrillo, Y. Huang, M. C. Santos, J. L. Vázquez. *Exponential convergence towards stationary states for the 1D porous medium equation with fractional pressure.* Journal of Differential Equations, 2015, 258(3): 736-763.

[8] A. Córdoba, D. Córdoba. *A maximum principle applied to quasi-geostrophic equations.* Communications in mathematical physics, 2004, 249(3): 511-528.

[9] N. Ju. *Existence and uniqueness of the solution to the dissipative 2D quasi-geostrophic equations in the Sobolev space.* Communications in Mathematical Physics, 2004, 251(2): 365-376.

[10] C. Miao, J. Wu, Z. Zhang. *Littlewood-Paley Theory and Applications to Fluid Dynamics Equations.* Monographson Modern Pure Mathematics, No. 142 (Science Press, Beijing, 2012).

[11] D. Stan, F. del Teso, J. L. Vázquez. *Transformations of self-similar solutions for porous medium equations of fractional type.* Nonlinear Analysis: Theory, Methods Applications, 2015, 119: 62-73.

[12] J. L. Vázquez. *Nonlinear diffusion with fractional Laplacian operators.* Nonlinear Partial Differential Equations. Springer Berlin Heidelberg, 2012: 271-298.

[13] X. Zhou, W. Xiao, J. Chen, *Fractional porous medium and mean field equations in Besov spaces*, Electron. J. Differential Equations, 2014 (2014), 1-14.

[1] Department of Information Technology, Nanjing Forest Police College, 210023 Nanjing, China
Email: zhouxuhuan@163.com

[2*] School of Applied Mathematics, Nanjing University of Finance and Economics, 210023 Nanjing, China
Email: xwltc123@163.com